

Peer Review of the Nickel Development Support Document

List of Key References

Agency for Toxic Substances and Disease Registry (ATSDR). 2005. Toxicological profile for Nickel. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service. Available online at: <http://www.atsdr.cdc.gov/toxprofiles/tp15.html>.

Circla AM, Bernabeo F, Ottoboni F, Ratti R. 1985. Nickel induced occupational asthma: Immunological and clinical aspects. IN: Progress in Nickel Toxicology. Eds. Brown S, Sunderman FW. International Union of Pure and Applied Chemistry, Blackwell Scientific Publications.

Doll R, Anderson A, Cooper WC, et al. 1990. Report of the International Committee on nickel carcinogenesis in man. *Scand J Work Environ Health* 16:1-82.

Enterline PE, Marsh GM. 1982. Mortality among workers in a nickel refinery and alloy manufacturing plant in West Virginia. *Journal of the National Cancer Institute*. 68(6):925-933.

Fernandez-Nieto M, Quirce S, Carnes J, Sastre J. 2006. Occupational asthma due to chromium and nickel salts. *Int Arch Occup Environ Health* 79:483-486.

Goodman JE, Prueitt RL, Dodge DG, Thakali S. 2009. Carcinogenicity assessment of water-soluble nickel compounds. Gradient Corporation. 20 University Rd, Cambridge, MA.

Graham JA, Miller FJ, Daniels MJ, et al. 1978. Influence of cadmium, nickel, and chromium on primary immunity in mice. *Environ Res* 16:77-87.

Grimsrud TK, Berge SR, Resmann F, et al. 2000. Assessment of historical exposures in a nickel refinery in Norway. *Scand J Work Environ Health* 26(4):338-345.

Grimsrud TK, Berge SR, Haldorsen T, Andersen A. 2002. Exposure to different forms of nickel and risk of lung cancer. *American Journal of Epidemiology* 156(12):1123-1132.

Grimsrud TK, Berge SR, Martinsen JI, Andersen A. 2003. Lung cancer incidence among Norwegian nickel-refinery workers 1953-2000. 2003. *JEM* 5:190-197.

Shihara Y, Kyono H, Serita F, Toya T, Kawashima H, Miyasaka M. 2002. Inflammatory responses and mucus secretion in rats with acute bronchiolitis induced by nickel chloride. *Inhalation Toxicology* 14:417-430.

Magnus K, Andersen A, Hogetveit AC. 1982. Cancer of respiratory organs among workers at a nickel refinery in Norway. *International Journal of Cancer* 30(6):681-685.

National Toxicology Program (NTP). 1996. NTP technical report on the toxicology and carcinogenesis studies of nickel sulfate hexahydrate (CAS no. 10101-97-0) in F344/N rats and B6C3F1s mice (inhalation studies). In NIH publication, National Toxicology Program technical report series; no. 454.

Seilkop SK, Oller AR. 2003. Respiratory cancer risks associated with low-level nickel exposure: an integrated assessment based on animal, epidemiological, and mechanistic data. *Regulatory Toxicology and Pharmacology* 37:173-190.

Serita F. 1999. An animal exposure system using ultrasonic nebulizer that generates well controlled aerosols from liquids. *Industrial Health* 37:82-87.

Texas Commission on Environmental Quality (TCEQ). 2006. Guidelines to develop effects screening levels, reference values, and unit risk factors. Chief Engineer's Office. RG-442: Texas Commission on Environmental Quality.

United States Environmental Protection Agency (U.S. EPA). 1986. Health assessment document for nickel and nickel compounds: final report. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development, Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office.